

THE MINIMAL VOLUME ORIENTABLE HYPERBOLIC 2-CUSPED 3-MANIFOLDS

IAN AGOL

ABSTRACT. We prove that the Whitehead link complement and the $(-2, 3, 8)$ pretzel link complement are the minimal volume orientable hyperbolic 3-manifolds with two cusps, with volume $3.66\dots = 4 \times \text{Catalan's constant}$. We use topological arguments to establish the existence of an essential surface which provides a lower bound on volume and strong constraints on the manifolds that realize that lower bound.

1. INTRODUCTION

Jorgensen and Thurston proved that the volumes of hyperbolic 3-manifolds are well-ordered. Moreover, if a volume is an n -fold limit point of smaller volumes (of order type ω^n), then there is a corresponding hyperbolic manifold of finite volume with precisely n orientable cusps. The smallest volume one-cusped orientable hyperbolic manifolds were identified by Cao and Meyerhoff [5], one of which is the figure-eight knot complement, with volume $2.0298\dots = 2V_3$, where $V_3 = 1.01494\dots$ is the volume of a regular ideal tetrahedron. Recently, Gabai, Meyerhoff and Milley have identified the smallest volume orientable hyperbolic 3-manifold to be the Fomenko-Matveev-Weeks manifold, with volume $0.9427\dots$ [11]. Moreover, they identify the 10 one-cusped orientable manifolds with volume < 2.848 (having five distinct volumes). It is an interesting question to find the smallest volume manifolds with n orientable cusps. Adams showed that an n -cusped hyperbolic manifold has volume $\geq nV_3$ [1]. For orientable manifolds with two cusps, this was improved by Yoshida to a lower bound of 2.43 in [22]. In this paper, we prove that the Whitehead link complement $\mathbb{W}$ and the $(-2, 3, 8)$ pretzel link complement $\mathbb{W}'$ are the minimal volume orientable hyperbolic 3-manifolds with two cusps, with volume $V_8 = 3.66\dots = 4 \times \text{Catalan's constant}$. These have been the smallest volume known 2-cusped orientable hyperbolic 3-manifolds for quite some time, and we take it as given that these are known to have this volume (see [21] or [20, p. 474]). In fact, in Theorem 3.5, we prove that if M is a hyperbolic manifold with a cusp and an essential surface disjoint from the cusp, then $\text{Vol}(M) \geq V_8$, from which the result for 2-cusped manifolds follows in Theorem 3.6 by a result of Culler and Shalen [7].

We now briefly outline the argument. Let M be a compact orientable manifold with at least one boundary component, whose interior $\text{int}(M)$ admits a finite volume hyperbolic metric and which contains a closed essential surface X disjoint from the boundary. If the

Received by the editors January 6, 2010.

2000 *Mathematics Subject Classification.* 57M.

Agol partially supported by NSF grant DMS-0504975, and the Guggenheim foundation.

complement of X has trivial “guts” (that is, the double $D(M \setminus X)$ of M split along X is a graph manifold), then we use the JSJ decomposition of the complement of X to show that the boundary may be used to “cut” up the surface X , repeating until we get a surface which has non-trivial guts. Essentially what we are doing is replacing the initial surface X which may have accidental parabolics, with a surface which has no accidental parabolics, but we must use a topological approach rather than geometric in order to keep track of the JSJ decomposition and get an acylindrical part of the guts in the end. We then apply a volume estimate of [2] and volume estimates of Miyamoto [13] giving lower bounds on volumes of hyperbolic manifolds with totally geodesic boundary, to get the volume lower bound on $\text{int}(M)$ (Theorem 3.5). A recent sharp estimate [3] implies that we may characterize completely the case of equality and identify the two smallest volume manifolds with two cusps by a simple combinatorial analysis (Theorem 3.6). The arguments of [2, 3] depend strongly on Perelman’s proof of the geometrization conjecture (see [19, 18, 12, 14, 6, 15]).

Acknowledgements: We thank Chris Atkinson and Jeff Weeks for helpful suggestions. We also thank Rupert Venzke and Danny Calegari for a helpful discussion about the minimal volume hyperbolic manifold with n cusps and for correcting a mistaken conjecture in the first version of the paper. We thank the referee for many helpful comments, and for pointing out a strengthening of the main theorem.

2. DEFINITIONS

In this section, we set up some notation and terminology for the theory of characteristic submanifolds, which is a relative version of the geometric decomposition.

For a surface $X \subset M$, we’ll use $M \setminus X$ to indicate the path-metric closure of $M \setminus X$. Let M be an irreducible, orientable manifold. We will say that a properly embedded surface is *essential* if it is π_1 -injective and $\partial\pi_1$ -injective. Let $(X, \partial X) \subset (M, \partial M)$ be a properly embedded essential surface. A *compressing annulus* for X is an embedding

$$i : (S^1 \times I, S^1 \times \{0\}, S^1 \times \{1\}) \hookrightarrow (M, X, \partial M)$$

such that

- i_* is an injection on π_1 ,
- $i(S^1 \times I) \cap X = i(S^1 \times \{0\})$, and
- $i(S^1 \times \{0\})$ is not isotopic in X to ∂X .


An *annular compression* of $(X, \partial X) \subset (M, \partial M)$ is a surgery along a compressing annulus $i : (S^1 \times I, S^1 \times \{0\}, S^1 \times \{1\}) \hookrightarrow (M, X, \partial M)$. Let U be a regular neighborhood of $i(S^1 \times I)$ in $M \setminus X$, let $\partial_1 U$ be the frontier of U in $M \setminus X$, and let $\partial_0 U = \partial U \cap (X \cup \partial M)$. Then let $X' = (X - \partial_0 U) \cup \partial_1 U$. The surface X' is the *annular compression* of X (see Figure 1). We remark that if X is essential, then X' is as well.

The notion of a *pared manifold* was defined by Thurston to give a topological characterization of geometrically finite hyperbolic 3-manifolds (see [16] or [4, Ch. 5]).

Definition 2.1. A *pared manifold* is a pair (M, P) where

- M is a compact, orientable irreducible 3-manifold and
- $P \subset \partial M$ is a union of essential annuli and tori in M ,

such that

FIGURE 1. An annular compression of a surface (cross the picture with S^1)

- every abelian, noncyclic subgroup of $\pi_1(M)$ is peripheral with respect to P (i.e., conjugate to a subgroup of the fundamental group of a component of P) and
- every map $\varphi : (S^1 \times I, S^1 \times \partial I) \rightarrow (M, P)$ that is injective on the fundamental groups deforms, as maps of pairs, into P .

P is called the *parabolic locus* of the **pared manifold** (M, P) . We denote by $\partial_0 M$ the surface $\partial M - \text{int}(P)$.

The motivation for the introduction of **pared manifolds** is the following geometrization theorem of Thurston:

Theorem 2.2. [4, Ch. 7] *If (M, P) is an oriented **pared 3-manifold** with nonempty boundary, then there exists a geometrically finite uniformization of (M, P) .*

The term uniformization means that there is a hyperbolic manifold N such that N is homeomorphic to $\text{int}(M)$, and whose parabolic subgroups correspond to P (see [4, Ch. 7] for more details).

Let (M, P) be a **pared manifold** such that $\partial_0 M$ is incompressible. There is a canonical set of essential annuli $(A, \partial A) \subset (M, \partial_0 M)$, called the *characteristic annuli*, such that $(P, \partial P) \subset (A, \partial A)$, and characterized (up to isotopy) by the property that they are the maximal collection of non-parallel essential annuli such that every other essential annulus $(B, \partial B) \subset (M, \partial_0 M)$ may be relatively isotoped to an annulus $(B', \partial B') \subset (M, \partial_0 M)$ so that $B' \cap A = \emptyset$. Each complementary component $L \subset M \setminus A$ is one of the following types:

- (1) $T^2 \times I$, a neighborhood of a torus component of P ,
- (2) $(S^1 \times D^2, S^1 \times D^2 \cap \partial_0 M)$, a solid torus with annuli in the boundary,
- (3) $(I - \text{bundles}, \partial I - \text{subbundles})$, where the I -bundles over the boundary are subsets of A , or
- (4) all essential annuli in $(L, \partial_0 M \cap L)$ are parallel in L into $(L \cap A, \partial(L \cap A))$.

The union of components of type (3), denoted $(W, \partial_0 W) \subset (M, \partial_0 M)$, is called the *window* of $(M, \partial_0 M)$. It is unique up to isotopy of pairs.

The **pared submanifold** $(M - W, \partial(M - W) - \partial_0 M)$ is denoted $Guts(M, P)$. Note that the parabolic locus of $Guts(M, P)$ will consist of characteristic annuli. If M is compact orientable and $\text{int}(M)$ admits a metric of finite volume, and $(X, \partial X) \subset (M, \partial M)$ is an essential surface, then define $Guts(X) = Guts(M \setminus X, \partial M \setminus \partial X)$. The components of type (4) are acylindrical **pared manifolds**, which have a complete hyperbolic structure of

finite volume with geodesic boundary [16]. We will let $\text{Vol}(Guts(M, P))$ denote the volume of this hyperbolic metric. If $D(M, P)$ is obtained by taking two copies of M and gluing them along the corresponding surfaces $\partial_0 M$, then $\text{Vol}(Guts(M, P)) = \frac{1}{2}\text{Vol}(D(M, P))$, where $\text{Vol}(D(M, P))$ is the simplicial volume of $D(M, P)$, *i.e.* the sum of the volumes of the hyperbolic pieces of the geometric decomposition. If $\text{Vol}(Guts(M, P)) = 0$, then M is a “book of I -bundles”, with “pages” consisting of W , and “spine” consisting of solid tori and $T^2 \times I$ (see [4, Example 2.10.4] for more information).

Let $V_8 = 4 \cdot K = 3.66\dots$, where K is Catalan’s constant

$$K = 1 - 1/9 + 1/25 - 1/49 + \dots + (-1)^n/(2n+1)^2 + \dots$$

Then V_8 is also the volume of a regular ideal octahedron in $\mathbb{H}^3$.

3. ESSENTIAL SURFACES

In this section, we prove the existence of an essential surface with non-trivial **guts** under certain hypotheses (Theorem 3.4). In the terminology of Culler and Shalen, we find a surface which is not a *fibroid* [9], *i.e.* whose complement is not a book of I -bundles. We start with an essential surface missing one cusp. If this surface has no accidental parabolics, we are done. If it does, we perform annular compressions until we get a surface with no accidental parabolics and which is not a fibroid.

Lemma 3.1. *Let (M, P) be a **pared manifold**, with **window** $(W, \partial_0 W) \subset (M, \partial_0 M)$. Suppose that there is a component $(J, K) \subset (W, \partial_0 W)$ which is an I -bundle such that $\chi(J) < 0$, and such that $K \cap P \neq \emptyset$. Let K_0 be a component of $K \cap P$. Then the surface $\partial_0 M \cup K_0$ is compressible in M .*

Proof. Let Q be a surface such that $I \rightarrow J \rightarrow Q$ is a fiber bundle. Then $I \rightarrow K \rightarrow \partial Q$ is a bundle over the boundary. Let $q_0 \subset \partial Q$ be the component of ∂Q such that K_0 fibers over q_0 . Let $(\alpha, \partial\alpha) \subset (Q, q_0)$ be an essential arc. Then there is a disk $D \subset J$ which fibers over α . The disk D is essential, which implies that $\partial_0 M \cup K_0$ is compressible. To see this, note that if D is non-separating in M , then it is essential. So we may assume that D separates M , and therefore D separates J . But J is π_1 -injective in M , and since α is essential in Q , D is essential in J . Thus, each component of $J - D$ is π_1 -injective in J , and therefore in M , and neither component of $J - D$ is simply-connected. Therefore neither component of $\overline{M - D}$ can be a ball, since it contains an non-simply connected π_1 -injective submanifold, which implies that D is essential, and therefore that $\partial_0 M \cup K_0$ is compressible. $\square$

Lemma 3.2. *Let M be an orientable compact manifold such that $\text{int}(M)$ is hyperbolic of finite volume and so that $P = \partial M$ is the pared locus of M . Let $X \subset M$ be an essential surface. Then $(M \setminus X, P \setminus \partial X)$ is a **pared manifold**.*

Proof. Since each component V of $M \setminus X$ is π_1 -injective, there is a covering space $\tilde{M} \rightarrow M$ such that there is a lift $M \setminus X \rightarrow \tilde{M}$ which is a homotopy equivalence. Any abelian non-cyclic subgroup of $\pi_1(V)$ must be peripheral in M and therefore in $\tilde{M}$ and $M \setminus X$. Any map $\varphi : (S^1 \times I, S^1 \times \partial I) \rightarrow (M \setminus X, P \setminus \partial X)$ that is injective on the fundamental groups deforms, as maps of pairs, into $P \setminus X \subset \tilde{M}$, since components of P correspond to cusps associated to a complete hyperbolic structure on $\text{int}(\tilde{M})$ induced from the hyperbolic

structure on $\text{int}(M)$. Thus, we see that $(M \setminus \setminus X, P \setminus \setminus \partial X)$ satisfies the hypotheses of a **pared manifold**. $\square$

Lemma 3.3. *Let M be an orientable compact manifold such that $\text{int}(M)$ is hyperbolic of finite volume, so that $P = \partial M$ is the pared locus of M . Let $X \subset M$ be an essential surface. If X has a compressing annulus, let X_1 be the surface obtained by performing an **annular compression** along X . Then X_1 has a pared annulus coming from ∂M in the boundary of one of its gut regions.*

Proof. Let $A \subset M \setminus \setminus X$ be a compressing annulus, with $\partial A = a_0 \cup a_1$, such that $a_0 \subset \partial M$, and $a_1 \subset X$ is a closed curve in X which is not boundary parallel in X . By Lemma 3.2, $(M \setminus \setminus X, P \setminus \setminus \partial X)$ is a **pared manifold**. Compression along A creates the essential surface X_1 (see Figure 1). There will be a new component of the pared locus of $M \setminus \setminus X_1$ which is an annulus with core a_0 . This annulus must be in a component of $\text{Guts}(X_1)$, because otherwise a_0 would lie in the boundary of the core of the **window** $W \subset M \setminus \setminus X_1$. This implies that X had a compression, since reversing the **annular compression** to obtain X from X_1 corresponds to adding a boundary annulus to an I -bundle of Euler characteristic < 0 , creating a compression by Lemma 3.1. $\square$

Theorem 3.4. *Let M be an orientable compact 3-manifold whose interior admits a hyperbolic metric of finite volume. Suppose that ∂M contains a torus T_1 , and that there is a 2-sided essential surface $X_0 \subset M$ such that $\partial X_0 \cap T_1 = \emptyset$. Then there is an essential surface $X \subset M$ such that $\chi(\text{Guts}(X)) < 0$.*

Proof. Let $\partial M = T_1 \cup \dots \cup T_n$, where T_i is a torus. Then $\partial X_0 \subset T_2 \cup \dots \cup T_n$. Suppose that $\chi(\text{Guts}(X_0)) = 0$. Then $M \setminus \setminus X_0$ is a book of I -bundles. Now, we perform a maximal sequence of **annular compressions** along ∂M to obtain a sequence of essential surfaces $X_1, X_2, \dots, X_k \subset M$, where X_{i+1} is obtained from X_i by an **annular compression**. Since X_0 is disjoint from T_1 , the component of $\text{Guts}(X_0)$ incident with T_1 must be of the form $T^2 \times I$, and thus X_0 has an **annular compression**. The compressing annulus comes from a curve in T^2 which is parallel to the boundary of a characteristic annulus of $M \setminus \setminus X_0$, and which is parallel to a curve in T_1 via a compressing annulus. For $i > 0$, X_i will be a 2-sided essential surface, with a component of the pared locus coming from ∂M lying in the boundary of a gut region of $M \setminus \setminus X_i$, by Lemma 3.3. If $\chi(\text{Guts}(X_i)) = 0$ for $i > 0$, then the components of $\text{Guts}(X_i)$ must be solid tori or tori $\times I$. Since M is simple, a pared annulus in ∂M intersect the boundary of a solid torus component of $\text{Guts}(X_i)$ must be primitive, and thus there is an **annular compression** of X_i (see Figure 2). Any region of $\text{Guts}(X_i)$ of the form $T^2 \times I$ will also give rise to an **annular compression** of X_i , as we observed for X_0 . Since $\chi(X_{i+1}) = \chi(X_i)$ and X_{i+1} has two more boundary components than X_i , the number of **annular compressions** is finite, and thus at some stage we arrive at X_k such that $\chi(\text{Guts}(X_k)) < 0$. Let $X = X_k$. $\square$

Theorem 3.5. *Let M be an orientable compact 3-manifold whose interior admits a hyperbolic metric of finite volume. Suppose that ∂M contains a torus T_1 , and that there is a 2-sided essential surface $X_0 \subset M$ such that $\partial X_0 \cap T_1 = \emptyset$. Then $\text{Vol}(M) \geq V_8$. If $\text{Vol}(M) = V_8$, then M has a decomposition into a single right-angled octahedron.*




FIGURE 2. An annular compression a of X_i gives X_{i+1} (cross the picture with S^1)

Proof. From Theorem 3.4, there is an essential surface $X \subset M$ such that $\chi(Guts(X)) \leq -1$. By [3, Theorem 5.5], we have $\text{Vol}(M) \geq \text{Vol}(Guts(X))$ (see also [2, Theorem 9.1] on which the method of proof of [3] is based). By [13, Theorem 4.2], $\text{Vol}(Guts(X)) \geq -V_8\chi(Guts(X))$. Thus, we have $\text{Vol}(M) \geq V_8$.

If $\text{Vol}(M) = V_8$, then $\text{Vol}(M) = \text{Vol}(Guts(X))$. We may assume that no complementary region of $M \setminus X$ is an I -bundle, since otherwise we could replace X with a surface which it double covers. By [3, Theorem 5.5], X is totally geodesic, and so $N = M \setminus X = Guts(X)$. Regardless of whether X is one- or two-sided, we have $\chi(X) = \frac{1}{2}\chi(\partial N) = \chi(N) = -1$, so X is either a punctured torus, a thrice-punctured sphere, or a twice-puncture $\mathbb{RP}^2$. Since N is an acylindrical manifold with $\chi(N) = -1$ and $\text{Vol}(N) = V_8$, by Miyamoto's theorem [13, Theorem 4.2] N must have a decomposition into a single regular octahedron of volume V_8 . Thus M decomposes into a single regular octahedron. $\square$

Remark: The referee indicated the previous theorem as a natural generalization of our main result, and pointed out that there is a sharp example (see [10, Section 6]). The example is m137 from the Snappea census [21], and is obtained by 0-framed surgery on a 2-component link L (see Figure 3). To see that m137 contains a closed essential surface, note that there is a thrice-punctured sphere P in the complement of m137 (see the grey surface in the second Figure 3). The surgery along the boundary slope of P gives $S^2 \times S^1$, since the other component of L is unknotted. However, since m137 is hyperbolic, it does not fiber over S^1 with fiber P , and therefore by [8, Theorem 2.0.3], m137 must contain a closed essential surface (which is obtained by carefully tubing together the boundary components of two copies of P).

Theorem 3.6. *Suppose that M is an orientable hyperbolic 3-manifold with two cusps, then $\text{Vol}(M) \geq V_8 = 3.66\dots$. If $\text{Vol}(M) = V_8$, then $M \cong \mathbb{W}$ or $M \cong \mathbb{W}'$.*



FIGURE 3. m137 realizes the lower bound in Theorem 3.5, and a thrice-punctured sphere P in m137

Proof. By [7, Theorem 3], there is an orientable connected essential separating surface $X_0 \subset M$ such that $\partial X_0 \cap (T_2 \cup \dots \cup T_n) = \emptyset$, and $\partial X_0 \cap T_1$ is a non-empty union of non-contractible simple closed curves. Moreover, we may assume that there is no compressing annulus of X_0 along T_1 . By Theorem 3.5, $\text{Vol}(M) \geq V_8$.

Now, suppose $\text{Vol}(M) = V_8$. By Theorem 3.5, M contains an essential surface X with $\chi(X) = -1$, and the complement of X has a decomposition into a regular octahedron.




FIGURE 4. A twice-punctured $\mathbb{RP}^2$ in the Whitehead and $(-2, 3, 8)$ -pretzel link complements

To finish the argument, we take an octahedron O and color the faces alternately black and white. The argument here is somewhat indirect. We know we have two examples, $\mathbb{W}, \mathbb{W}'$, and we need to show that there are no more. We glue the black faces of O together in all possible ways to get the possible manifolds N . Up to isometry, we get four manifolds $N_i, i = 1, 2, 3, 4$ with totally geodesic boundary and rank one cusps (see Figure 5, where the black faces being paired are labelled A and B). Then ∂N_1 is two pairs of pants, and ∂N_i is a single four-punctured sphere, for $i = 2, 3, 4$ (∂N_i is shown on the right of Figure 5).

Gluing the two boundary components of N_1 together in a way that gives two cusps gives $\mathbb{W}$ the Whitehead link complement. There are three other manifolds which have boundary a four-punctured sphere. The boundary gets glued to itself by an antipodal map. To get a manifold with two cusps from N_i , the antipodal map must also identify the rank one cusps of ∂N_i which share a common rank one cusp of N_i . Only N_2 and N_4 have boundary which admits an antipodal isometry, so that the quotient is $X \cong \mathbb{RP}^2 - \{x, y\}$. To see that ∂N_3 cannot have such an antipodal map, notice that ∂N_3 has isometry group S_4 , for which there is no isometry which acts as an antipodal map. The antipodal isometry preserves the induced triangulation of ∂N_i shown on the right of Figure 5 for $i = 2, 4$. One may check that the quotient by these isometries give a manifold with two cusps. Any other isometry which had this property must be the same, since the product of two such isometries must fix all four cusps of ∂N_i , which implies that it is the identity since this is the only holomorphic map of S^2 fixing four points. One quotient by this map gives back $\mathbb{W}$, since $\mathbb{W}$ has a twice-punctured $\mathbb{RP}^2$ in its complement (see Figure 4). The other example must give the $(-2, 3, 8)$ -pretzel link complement $\mathbb{W}'$ (in fact, N_2 corresponds to $\mathbb{W}'$ and N_4 corresponds to $\mathbb{W}$, after one quotients by the unique antipodal map of the boundaries). $\square$

4. CONCLUSION

It would be interesting to find the minimal volume orientable hyperbolic manifolds with n cusps. We conjecture that the minimal volume n -cusped manifold is realized by a hyperbolic chain link which is minimally “twisted” (see [17] for the definition of chain links) for $n \leq 10$. Rupert Venzke has pointed out to us that for links with n cusps, where $n \geq 11$, the $n - 1$ -fold cyclic cover $\mathbb{W}_n$ over one component of $\mathbb{W}$ has volume less than the smallest volume twist link with n cusps, which is a Dehn filling on $\mathbb{W}_{n+1}$. It is probably impossible to use the methods in this paper to determine the smallest volume hyperbolic manifold with n cusps. However, it should be possible to prove the following:

Conjecture 4.1. Let v_n be the minimal volume of an n -cusped orientable hyperbolic 3-manifold. Then

$$\lim_{n \rightarrow \infty} v_n/n = V_8.$$

As mentioned before, Adams has shown that $v_n/n \geq V_3$ [1]. It may also be possible to prove:

Conjecture 4.2. The minimal limit volume of non-orientable hyperbolic 3-manifolds is V_8 .

In the Snappea census of 3-manifolds [21], there are four manifolds of volume V_8 which are non-orientable with a single orientable cusp, which would correspond to the smallest limit volumes of non-orientable manifolds if this conjecture were true. It seems possible that one could prove this conjecture by proving the existence of an essential surface disjoint from the orientable cusp with non-trivial guts.

REFERENCES

- [1] Colin C. Adams, *Volumes of N -cusped hyperbolic 3-manifolds*, J. London Math. Soc. (2) **38** (1988), no. 3, 555–565.

- [2] Ian Agol, Peter A. Storm, and William P. Thurston, *Lower bounds on volumes of hyperbolic Haken 3-manifolds*, J. Amer. Math. Soc. **20** (2007), no. 4, 1053–1077 (electronic), arXiv:math.DG/0506338, With an appendix by Nathan Dunfield.
- [3] Danny Calegari, Michael H. Freedman, and Kevin Walker, *Positivity of the universal pairing in 3 dimensions*, preprint, March 2008, arXiv:0802.3208.
- [4] Richard D. Canary and Darryl McCullough, *Homotopy equivalences of 3-manifolds and deformation theory of Kleinian groups*, Mem. Amer. Math. Soc. **172** (2004), no. 812, xii+218.
- [5] Chun Cao and G. Robert Meyerhoff, *The orientable cusped hyperbolic 3-manifolds of minimum volume*, Invent. Math. **146** (2001), no. 3, 451–478.
- [6] Huai-Dong Cao and Xi-Ping Zhu, *A complete proof of the Poincaré and geometrization conjectures—application of the Hamilton-Perelman theory of the Ricci flow*, Asian J. Math. **10** (2006), no. 2, 165–492, arXiv:math/0612069v1.
- [7] M. Culler and P. B. Shalen, *Bounded, separating, incompressible surfaces in knot manifolds*, Invent. Math. **75** (1984), no. 3, 537–545.
- [8] Marc Culler, C. McA. Gordon, J. Luecke, and Peter B. Shalen, *Dehn surgery on knots*, Ann. of Math. (2) **125** (1987), no. 2, 237–300.
- [9] Marc Culler and Peter B. Shalen, *Volumes of hyperbolic Haken manifolds. i.*, Inventiones Mathematicae **118** (1994), no. 2, 285–329.
- [10] Nathan M. Dunfield, *Examples of non-trivial roots of unity at ideal points of hyperbolic 3-manifolds*, Topology **38** (1999), no. 2, 457–465.
- [11] David Gabai, Robert Meyerhoff, and Peter Milley, *Minimum volume cusped hyperbolic three-manifolds*, J. Amer. Math. Soc. **22** (2009), no. 4, 1157–1215, arXiv:0705.4325.
- [12] Bruce Kleiner and John Lott, *Notes on Perelman’s papers*, Geom. Topol. **12** (2008), no. 5, 2587–2855, arXiv:math.DG/0605667.
- [13] Yosuke Miyamoto, *Volumes of hyperbolic manifolds with geodesic boundary*, Topology **33** (1994), no. 4, 613–629.
- [14] John Morgan and Gang Tian, *Ricci flow and the Poincaré conjecture*, Clay Mathematics Monographs, vol. 3, American Mathematical Society, Providence, RI, 2007, arXiv:math/0607607v2.
- [15] John Morgan and Gang Tian, *Completion of the proof of the geometrization conjecture*, preprint, 2008, arXiv.org:0809.4040.
- [16] John W. Morgan, *On Thurston’s uniformization theorem for three-dimensional manifolds*, The Smith conjecture (New York, 1979), Academic Press, Orlando, FL, 1984, pp. 37–125.
- [17] Walter D. Neumann and Alan W. Reid, *Arithmetic of hyperbolic manifolds*, Topology ’90 (Columbus, OH, 1990), Ohio State Univ. Math. Res. Inst. Publ., vol. 1, de Gruyter, Berlin, 1992, pp. 273–310.
- [18] Grisha Perelman, *Ricci flow with surgery on three-manifolds*, arXiv:math.DG/0303109.
- [19] ———, *The entropy formula for the Ricci flow and its geometric applications*, arXiv:math.DG/0211159.
- [20] John G. Ratcliffe, *Foundations of hyperbolic manifolds*, second ed., Graduate Texts in Mathematics, vol. 149, Springer, New York, 2006.
- [21] Jeffery Weeks, *SnapPea: A computer program for creating and studying hyperbolic 3-manifolds*, www.geometrygames.org.
- [22] Han Yoshida, *Volumes of orientable 2-cusped hyperbolic 3-manifolds*, Kobe J. Math. **18** (2001), no. 2, 147–161.

UNIVERSITY OF CALIFORNIA, BERKELEY, 970 EVANS HALL #3840, BERKELEY, CA 94720-3840
 E-mail address: ianagol@math.berkeley.edu




FIGURE 5. Gluing faces A and B of an octahedron together in pairs to get manifolds with geodesic boundary of volume V_8